


IN THE CLAIMS:

Claim 1 (currently amended and reformatted): A pyrimidine derivative compound of
the Formula I




I

wherein

m is 0, 1, 2 or 3 and

each R¹ group, which may be the same or different, is selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, sulphamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula :



wherein X¹ is a direct bond or is selected from O, S, SO, SO₂, N(R⁴), CO, CH(OR⁴), CON(R⁴), N(R⁴)CO, SO₂N(R⁴), N(R⁴)SO₂, OC(R⁴)₂, SC(R⁴)₂ and N(R⁴)C(R⁴)₂, wherein each R⁴ is hydrogen or (1-6C)alkyl, and Q² is aryl-(1-6C)alkyl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, or (R¹)_m is (1-3C)alkylenedioxy, and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R¹ substituent is optionally separated by the insertion of a group selected from O,

S, SO, SO₂, N(R⁵), CO, CH(OR⁵), CON(R⁵), N(R⁵)CO, SO₂N(R⁵) and N(R⁵)SO₂

wherein R⁵ is hydrogen or (1-6C)alkyl,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R¹

optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-

6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,


N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-

6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-

6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-

6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or

from a group of the formula :



wherein X² is a direct bond or is selected from O and N(R⁷), wherein R⁷ is

hydrogen or (1-6C)alkyl, and Q³ is aryl, aryl-(1-6C)alkyl, heteroaryl,

heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and any Q³

group optionally bears 1 or 2 substituents, which may be the same or different,

selected from halogeno, trifluoromethyl, cyano, hydroxy, amino,

(1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2

oxo or thioxo substituents,

and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each

said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a

substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-

6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-

6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-

6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-

6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-

(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-


6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino;

a' R³ is hydrogen, halogeno or (1-6C)alkyl;

n is 0, 1 or 2 and each R² group, which may be the same or different, is selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, (1-6C)alkoxycarbonyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

p is 0, 1, 2, 3 or 4; and

Q¹ is aryl or heteroaryl and Q¹ is optionally substituted with 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino or with a (1-3C)alkylenedioxy group, or from a group of the formula :



wherein X³ is a direct bond or is selected from O and N(R⁸), wherein R⁸ is hydrogen or (1-6C)alkyl, and Q⁴ is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and any Q⁴ group optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino,

(1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino,

and wherein any heterocyclyl group within a substituent on Q¹ optionally bears 1 or 2 oxo or thioxo substituents,

and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a Q¹ substituent is optionally separated by the insertion of a group selected from O, S, SO, SO₂, N(R⁹), CO, CH(OR⁹), CON(R⁹), N(R⁹)CO, SO₂N(R⁹) and N(R⁹)SO₂ wherein R⁹ is hydrogen or (1-6C)alkyl,


and wherein any CH₂ or CH₃ group within a Q¹ group optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a

a'

substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino; or a pharmaceutically-acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 2 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 0, 1, 2 or 3, and each R¹ group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylthio, methylsulphinyl, methylsulphonyl, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, propargylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl and acetamido, or from a group of the formula :



wherein X¹ is a direct bond or is selected from O, NH and N(Me) and Q² is benzyl, 2-furylmethyl, 3-furylmethyl, 2-thienylmethyl, 3-thienylmethyl, 1-imidazolylmethyl, 2-imidazolylmethyl, 2-imidazol-1-ylethyl, 3-imidazol-1-ylpropyl, 4-imidazol-1-ylbutyl, 2-oxazolylmethyl, 4-oxazolylmethyl, 5-oxazolylmethyl, 2-thiazolylmethyl, 4-thiazolylmethyl, 5-thiazolylmethyl, 1,2,3-triazol-1-ylmethyl, 2-(1,2,3-triazol-1-yl)ethyl, 3-(1,2,3-triazol-1-yl)propyl, 1,2,4-triazol-1-ylmethyl, 2-(1,2,4-triazol-1-yl)ethyl, 3-(1,2,4-triazol-1-yl)propyl, 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 2-pyrid-2-ylethyl, 2-pyrid-3-ylethyl, 2-pyrid-4-ylethyl, 3-pyrid-2-ylpropyl,

Q1
3-pyrid-3-ylpropyl, 3-pyrid-4-ylpropyl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl, homopiperidin-1-yl, homopiperidin-2-yl, homopiperidin-3-yl, homopiperidin-4-yl, piperazin-1-yl, homopiperazin-1-yl, pyrrolidin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-1-ylpropyl, pyrrolidin-2-ylmethyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-2-ylpropyl, pyrrolidin-3-ylmethyl, 2-pyrrolidin-3-ylethyl, 3-pyrrolidin-3-ylpropyl, imidazolidin-1-ylmethyl, 2-imidazolidin-1-ylethyl, 3-imidazolidin-1-ylpropyl, imidazolidin-2-ylmethyl, 2-imidazolidin-2-ylethyl, 3-imidazolidin-2-ylpropyl, morpholinomethyl, 2-morpholinoethyl, 3-morpholinopropyl, morpholin-2-ylmethyl, 2-morpholin-2-ylethyl, 3-morpholin-2-ylpropyl, morpholin-3-ylmethyl, 2-morpholin-3-ylethyl, 3-morpholin-3-ylpropyl, tetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(tetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(tetrahydro-4H-1,4-thiazin-4-yl)propyl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-ylmethyl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, piperidinomethyl, 2-piperidinoethyl, 3-piperidinopropyl, piperidin-2-ylmethyl, 2-piperidin-2-ylethyl, 3-piperidin-2-ylpropyl, piperidin-3-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-3-ylpropyl, piperidin-4-ylmethyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, homopiperidin-1-ylmethyl, 2-homopiperidin-1-ylethyl, 3-homopiperidin-1-ylpropyl, homopiperidin-2-ylmethyl, 2-homopiperidin-2-ylethyl, 3-homopiperidin-2-ylpropyl, homopiperidin-3-ylmethyl, 2-homopiperidin-3-ylethyl, 3-homopiperidin-3-ylpropyl, homopiperidin-4-ylmethyl, 2-homopiperidin-4-ylethyl, 3-homopiperidin-4-ylpropyl, piperazin-1-ylmethyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, piperazin-2-ylmethyl, 2-piperazin-2-ylethyl, 3-piperazin-2-ylpropyl, homopiperazin-1-ylmethyl, 2-homopiperazin-1-ylethyl, 3-homopiperazin-1-ylpropyl, homopiperazin-2-ylmethyl, 2-homopiperazin-2-ylethyl or 3-homopiperazin-2-ylpropyl,

a¹ and wherein a single pair of adjacent carbon atoms in a (2-6C)alkylene chain within a R¹ substituent is optionally separated by the insertion of a group selected from O and NH,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R¹ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,


and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents,

and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and diethylamino;

each of n and p is 0;

R³ is hydrogen or methyl; and

Q¹ is phenyl, 2-furyl, 2-thienyl, 4-oxazolyl, 5-isoxazolyl, 2- or 4-imidazolyl, 3- or 4-pyrazolyl, 4-thiazolyl, 5-isothiazolyl, 2-, 3- or 4-pyridyl, 4-pyridazinyl, 4- or 5-pyrimidinyl, 2- or 6-benzofuranyl, 2- or 6-indolyl, 2- or 6-benzothiophenyl, 2- or 6-quinolyl or 2- or 4-dibenzofuranyl which optionally bears 1, 2 or 3 substituents, , which may be the same or different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, acetamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino, 1-azetidiny, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, 1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl, 1-piperidinyl, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or from a group of the formula :



a' wherein X³ is a direct bond or is selected from O, NH and N(Me) and Q⁴ is phenyl, 2- or 3-furyl, 2- or 3-thienyl, 1- or 2-imidazolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 2-, 3- or 4-pyridyl, 3- or 4-pyridazinyl, 2- or 4-pyrimidinyl or 2-pyrazinyl, and any Q⁴ group optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, bromo, trifluoromethyl, hydroxy, amino, methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino,

and wherein any heterocyclyl group within Q¹ optionally bears 1 or 2 oxo substituents,

and wherein any CH₂ or CH₃ group within a Q¹ group optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino;

or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 3 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 0, 1 or 2 and each R¹ group, which may be the same or different, is selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, amino, carbamoyl, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, N-methylcarbamoyl, N,N-dimethylcarbamoyl, acetamido, benzyloxy, benzylamino, N-benzyl-N-methylamino, 2-furylmethoxy, 3-furylmethoxy, 2-imidazol-1-ylethylamino, 3-imidazol-1-ylpropylamino, 2-(1,2,4-triazol-1-yl)ethylamino, 3-(1,2,4-triazol-1-yl)propylamino, 2-pyridylmethoxy, 3-pyridylmethoxy, 4-pyridylmethoxy, pyrrolidin-1-yl, pyrrolidin-3-yloxy, pyrrolidin-3-ylamino, N-methyl-N-(3-pyrrolidinyl)amino, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino, N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy,

a'
homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino, pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy, pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethoxy, 3-imidazolidin-1-ylpropoxy, imidazolidin-2-ylmethoxy, 2-imidazolidin-2-ylethoxy, 3-imidazolidin-2-ylpropoxy, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-imidazolidin-2-ylethylamino, 3-imidazolidin-2-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-morpholinoethylamino, 3-morpholinopropylamino, morpholin-2-ylmethoxy, 2-morpholin-2-ylethoxy, 3-morpholin-2-ylpropoxy, 2-morpholin-2-ylethylamino, 3-morpholin-2-ylpropylamino, morpholin-3-ylmethoxy, 2-morpholin-3-ylethoxy, 3-morpholin-3-ylpropoxy, 2-morpholin-3-ylethylamino, 3-morpholin-3-ylpropylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino, piperidin-2-ylmethoxy, 2-piperidin-2-ylethoxy, 3-piperidin-2-ylpropoxy, piperidin-2-ylmethylamino, 2-piperidin-2-ylethylamino, 3-piperidin-2-ylpropylamino, piperidin-3-ylmethoxy, 2-piperidin-3-ylethoxy, 3-piperidin-3-ylpropoxy, piperidin-3-ylmethylamino, 2-piperidin-3-ylethylamino, 3-piperidin-3-ylpropylamino, piperidin-4-ylmethoxy, 2-piperidin-4-ylethoxy, 3-piperidin-4-ylpropoxy, piperidin-4-ylmethylamino, 2-piperidin-4-ylethylamino, 3-piperidin-4-ylpropylamino, 2-homopiperidin-1-ylethoxy, 3-homopiperidin-1-ylpropoxy, 2-homopiperidin-1-ylethylamino, 3-homopiperidin-1-ylpropylamino, homopiperidin-2-ylmethoxy, homopiperidin-2-ylmethylamino, homopiperidin-3-ylmethoxy, homopiperidin-3-ylmethylamino, homopiperidin-4-ylmethoxy, homopiperidin-4-ylmethylamino, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino, piperazin-2-ylmethoxy, piperazin-2-ylmethylamino, 2-piperazin-2-ylethoxy, 3-piperazin-2-ylpropoxy, 2-

Q' piperazin-2-ylethylamino, 3-piperazin-2-ylpropylamino, 2-homopiperazin-1-ylethoxy, 3-homopiperazin-1-ylpropoxy, 2-homopiperazin-1-ylethylamino, 3-homopiperazin-1-ylpropylamino, homopiperazin-2-ylmethoxy or homopiperazin-2-ylmethylamino,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R¹ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, trifluoromethyl, amino, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy propoxy, isopropoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents,

and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylsulphonyl, methylamino, ethylamino, dimethylamino and diethylamino;

each of n and p is 0;

R³ is methyl; and

Q¹ is phenyl or 3- or 4-pyridyl which optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, bromo, trifluoromethyl, cyano, amino, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, dipropylamino, N-ethyl-N-methylamino, N-methyl-N-propylamino, acetamido, N-methylacetamido, methanesulphonylamino, ethanesulphonylamino, N-methylmethanesulphonylamino, 1-azetidiny, 3-pyrrolin-1-yl, 1-pyrrolidinyl, morpholino, 1-piperidinyl, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, and wherein any CH₂ or CH₃ group within a Q¹ group optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino;

or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Q' Claim 4 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 0, 1 or 2 and each R¹ group, which may be the same or different, is selected from fluoro, chloro, bromo, amino, carbamoyl, methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, pyrrolidin-3-yloxy, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino, N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy, homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino, pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy, pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-morpholinoethylamino, 3-morpholinopropylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino or piperazin-2-ylmethoxy, and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 substituents, which may be the same or different, selected from hydroxy, fluoro, chloro, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl, and wherein any heterocyclyl group within a substituent on R¹ optionally bears an oxo substituent,

Q¹ and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methylamino, ethylamino, dimethylamino and diethylamino; each of n and p is 0; R³ is methyl; and Q¹ is phenyl or 4-pyridyl which optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, amino, methyl, methoxy, methylamino, ethylamino, dimethylamino, diethylamino, 1-pyrrolidinyl, morpholino, piperidino, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, and wherein any CH₂ or CH₃ group within a Q¹ group optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methoxy, ethoxy, methylamino, ethylamino, dimethylamino and diethylamino; or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 5 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 1 or 2 and each R¹ group, which may be the same or different, is selected from chloro, carbamoyl, 3-dimethylaminopropoxy, 3-dimethylamino-2,2-dimethylpropoxy, methylthio, 3-diethylaminopropylamino, 3-dimethylamino-2,2-dimethylpropylamino, 3-dimethylamino-2-hydroxypropylamino, N-isopropylpyrrolidin-3-yloxy, piperidin-4-yloxy, N-methylpiperidin-4-yloxy, N-ethylpiperidin-3-ylamino, N-methylpiperidin-4-ylamino, N-methyl-N-(N-methylpiperidin-4-yl)amino, 3-pyrrolidin-1-ylpropylamino, N-methylpyrrolidin-2-ylmethoxy, 2-(N-methylpyrrolidin-2-yl)ethoxy, 2-(N-methylpyrrolidin-2-yl)ethylamino, N-methylpiperidin-3-ylmethoxy and N,N'-dimethylpiperazin-2-ylmethoxy;

each of n and p is 0;

R³ is methyl; and

Q¹ is phenyl which bears 1 or 2 substituents, which may be the same or different, selected from fluoro, trifluoromethyl, dimethylamino, 1-pyrrolidinyl, morpholino, piperidino,

a' 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or Q¹ is 4-pyridyl which bears 1 substituent selected from dimethylamino, 1-pyrrolidinyl, morpholino, piperidino, 1-homopiperidinyl, 1-piperazinyl and 1-homopiperazinyl, or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 6 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 0, 1 or 2 and each R¹ group, which may be the same or different, is selected from fluoro, chloro, bromo, amino, carbamoyl, methoxy, ethoxy, propoxy, methylthio, methylamino, ethylamino, propylamino, isopropylamino, butylamino, allylamino, dimethylamino, diethylamino, dipropylamino, N-allyl-N-methylamino, pyrrolidin-3-yloxy, morpholino, tetrahydro-4H-1,4-thiazin-4-yl, piperidino, piperidin-3-yloxy, piperidin-4-yloxy, piperidin-3-ylamino, piperidin-4-ylamino, N-methyl-N-(3-piperidinyl)amino, N-methyl-N-(4-piperidinyl)amino, homopiperidin-1-yl, homopiperidin-3-yloxy, homopiperidin-4-yloxy, piperazin-1-yl, homopiperazin-1-yl, 2-pyrrolidin-1-ylethoxy, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethylamino, 3-pyrrolidin-1-ylpropylamino, pyrrolidin-2-ylmethoxy, 2-pyrrolidin-2-ylethoxy, 3-pyrrolidin-2-ylpropoxy, pyrrolidin-2-ylmethylamino, 2-pyrrolidin-2-ylethylamino, 3-pyrrolidin-2-ylpropylamino, pyrrolidin-3-ylmethoxy, 2-pyrrolidin-3-ylethoxy, 3-pyrrolidin-3-ylpropoxy, pyrrolidin-3-ylmethylamino, 2-pyrrolidin-3-ylethylamino, 3-pyrrolidin-3-ylpropylamino, 2-imidazolidin-1-ylethylamino, 3-imidazolidin-1-ylpropylamino, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-morpholinoethylamino, 3-morpholinopropylamino, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-piperidinoethylamino, 3-piperidinopropylamino, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, 2-piperazin-1-ylethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethylamino, 3-piperazin-1-ylpropylamino or piperazin-2-ylmethoxy, and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 substituents, which may be the same or different, selected from hydroxy, fluoro,

a' chloro, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, methoxycarbonyl, ethoxycarbonyl and tert-butoxycarbonyl,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears an oxo substituent,

and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group 1 or 2 methyl substituents or a substituent selected from hydroxy, amino, methylamino, ethylamino, dimethylamino and diethylamino;

each of n and p is 0;

R³ is methyl; and

Q¹ is 4-dibenzofuranyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl, amino, methyl, methoxy, methylamino, ethylamino, dimethylamino and diethylamino;

or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 7 (currently amended and reformatted): A pyrimidine compound ~~derivative of~~ the Formula I according to claim 1 wherein

m is 1 and the R¹ group is selected from chloro, carbamoyl, methoxy, ethoxy,

3-dimethylaminopropoxy, 3-dimethylamino-2,2-dimethylpropoxy, methylthio,

N-isopropylpyrrolidin-3-yloxy, piperidin-4-yloxy, N-methylpiperidin-4-yloxy,

N-ethylpiperidin-4-yloxy, N-propylpiperidin-4-yloxy, N-methylpyrrolidin-2-ylmethoxy,

2-(N-methylpyrrolidin-2-yl)ethoxy, N-methylpiperidin-3-ylmethoxy and

N,N'-dimethylpiperazin-2-ylmethoxy;

each of n and p is 0;

R³ is methyl; and

Q¹ is 4-dibenzofuranyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from fluoro, chloro, trifluoromethyl and dimethylamino,


or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.

Claim 8 (currently amended and reformatted): A pyrimidine compound ~~derivative~~ of the Formula I according to claim 1 selected from :-

a' 6-carbamoyl-2-chloro-4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]pyrimidine,
4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]-2-methylthiopyrimidine,
2-chloro-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine,
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(N-methylpiperidin-4-yloxy)pyrimidine,
2-(3-dimethylaminopropoxy)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine,
2-(3-dimethylamino-2,2-dimethylpropoxy)-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine,
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-(N-methylpiperidin-3-ylmethoxy)pyrimidine,
2-[N-methyl-N-(N-methylpiperidin-4-yl)amino]-4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]pyrimidine,
4-[2-methyl-5-(2-morpholinopyrid-4-ylcarbonylamino)anilino]-2-[2-(N-methylpyrrolidin-2-yl)ethylamino]pyrimidine,
2-(3-dimethylamino-2,2-dimethylpropoxy)-4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]pyrimidine,
4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]-2-(N-methylpiperidin-4-yloxy)pyrimidine,
4-[5-(3-fluoro-5-morpholinobenzamido)-2-methylanilino]-2-(N-propylpiperidin-4-yloxy)pyrimidine,
4-[5-(4-dibenzofuranylcarbonylamino)-2-methylanilino]-2-(N-methylpiperidin-4-yloxy)pyrimidine and
4-[5-(4-dibenzofuranylcarbonylamino)-2-methylanilino]-2-(3-dimethylamino-2,2-dimethylpropoxy)pyrimidine;
or a pharmaceutically acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof.


a' Claim 9 (currently amended and reformatted): A process for preparing aA pyrimidine compound derivative of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof, according to claim 1 may be prepared by the process of:-

(a) reacting an aniline of the Formula II




II

with an acid of the Formula III, or a reactive derivative thereof,




under standard amide bond forming conditions, wherein variable groups are as defined in claim 1 and wherein any functional groups are protected if necessary;

(b) the reaction of an activated heteroaryl compound of the Formula V



V

wherein L is a displaceable group, with an aniline of the Formula VII



VII

wherein variable groups are as defined in claim 1 and wherein any functional group is protected if necessary;

(c) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein R¹ or a substituent on Q¹ is an amino, (1-6C)alkylamino,

a' di-[(1-6C)alkyl]amino, substituted (1-6C)alkylamino, substituted di-[(1-6C)alkyl]amino, a N-linked heterocyclyl substituent or a heterocyclylamino substituent, the reaction of an appropriate amine with a pyrimidine compound ~~derivative~~ of the Formula I wherein R¹ or a substituent on Q¹ as appropriate is a suitable displaceable group and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary;

- (d) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein R¹ or a substituent on Q¹ is a (1-6C)alkoxy or substituted (1-6C)alkoxy substituent or a heterocyclyloxy substituent, the reaction of an appropriate alcohol with a pyrimidine compound ~~derivative~~ of the Formula I wherein R¹ or a substituent on Q¹ as appropriate is a suitable displaceable group and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary;
- (e) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein m is 0, the cleavage of a compound of the Formula I, wherein m is 1, 2 or 3 and each R¹ substituent is a halogeno group and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary;
- (f) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein R¹ or Q¹ contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, the alkylation of a pyrimidine compound ~~derivative~~ of the Formula I wherein R¹ or Q¹ contains a hydroxy group or a primary or secondary amino group as appropriate, and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary;
- (g) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein R¹ is a hydroxy group, the cleavage of a compound of the Formula I, wherein R¹ is a halogeno group and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary; or
- (h) for a compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester thereof, wherein R¹ is a (1-6C)alkylsulphinyl or (1-6C)alkylsulphonyl

Q' group, the oxidation of a compound of the Formula I, wherein R¹ is a (1-6C)alkylthio group and wherein other variable groups are as defined in claim 1 and wherein any functional group is protected if necessary,

and thereafter

- (i) removing any protecting groups; and
- (ii) optionally forming a pharmaceutically-acceptable salt or in-vivo-cleavable ester.

Claim 10 (currently amended): A pharmaceutical composition which comprises a pyrimidine compound ~~derivative~~ of the Formula I, or a pharmaceutically-acceptable or in-vivo-cleavable ester formed from an available carboxy group thereof, as defined in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.

Claims 11-12 (cancelled).

Claim 13 (new): A method of a treating disease or medical condition mediated by the production or effect of p38 kinase which comprises administering to a warm-blooded animal in need thereof a p38 kinase inhibiting amount of a pyrimidine compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof, as defined in claim 1.

Claim 14 (new): A method of a treating disease or medical condition mediated by the production or effect of TNF or IL-1 which comprises administering to a warm-blooded animal in need thereof a TNF or IL-1 inhibiting amount of a pyrimidine compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof, as defined in claim 1.

Claim 15 (new): A method for inhibiting the production or effect of TNF or IL-1 in a warm-blooded animal in need thereof comprising administering to said animal a TNF or IL-1 inhibiting amount of a pyrimidine compound of the Formula I, or a pharmaceutically-acceptable salt or in-vivo-cleavable ester formed from an available carboxy group thereof, as defined in claim 1.